

Practice Guidelines

For Family Physicians
Volume 6

Preface

Primary Health Care, PHC (through the Family Medicine) is the cornerstone to achieve the Universal Health Coverage which is considered as the safety net for implementing the Social Health Insurance.

The Ministry of Health and Population aims at improving the quality of health care service provided for all population groups in Egypt especially at Primary Health Care level. These goals couldn't be achieved without the development and capacity building of human resources for of health focusing on PHC Doctors (Family Physicians).

Family Practice is the core business in providing PHC services, so, developing and updating the Practice Guideline and Treatment Protocol is the foundation of offering quality care.

Through applying these Guideline the Family Physicians are capable and committed to the milestones and benchmarks of providing the basic benefits package of PHC services with the agreed and standard level of quality.

It's also mandatory to monitor and supervise the performance of providers and satisfaction of clients through parallel and comprehensive tools.

We are sure that Egyptian PHC Doctors can perfectly maximize the befits of utilizing these Guidelines for improving health of our population and subsequently the quality of life that leads the sustainable development for our Nation.

H.E. Prof. Dr.

Ahmed Emad El Din Rady

Minister of Health and Population

Egypt

This new edition of "Practice Guidelines For Family Physician" was updated, reprinted and issued on April, 2016 by the Ministry of Health and Population, Egypt

Message from His Excellency

Prof. Dr. Hatem El Gabaly

Comprehensive development and modernization is one of Egypt's priorities and pursued objectives. Out of this rule, we are committed towards improving the quality of health care services available for all Egyptians; adults, children, the poor and the well-off.

The Ministry of Health and Population has adopted, as a top priority, developing current systems to provide and finance health services in guidance and vision of the political leadership to ensure high quality in service provision and meet needs and expectations of the population as well as keeping up with top-notch developments at all levels: Primary, Secondary and Tertiary.

This vision has been translated into a promising and ambitious Five Years Plan to institutionalize the Health Sector Reform Program on the national level. The plan is focusing on implementing the Family Health Model at all primary health care facilities in the 27 Governorates.

Our dream has been realized into a competent program of Health Sector Reform aiming to provide every person with high quality health services. These include physical, psychological and social welfare, which translate into high production and progress for our cherished Country, Egypt.

I am delighted to introduce to one of the important publications for the Sector of Technical Support and Projects, representing a great team effort "The Practice Guidelines for Family Physicians" for the family physician at all Family Health Units of MOHP Distributed all over the Country.

Prof. Dr. Hatem El Gabaly

Minister of Health and Population

Preface

The Ministry of Health and population is working diligently to achieve equal and available quality health services for all citizens of Egypt. Our objective is to shape national policies for the goal of advancing health care delivery in all parts of the country.

Six years ago, the Ministry has adopted new policies and strategies in order to provide basic health services of high quality for all citizens in the framework of the Family Health Model. This has led to introducing new financing mechanisms that ensure the sustainability of finance and resources, and availability of affordable services along with effectiveness and efficiency of these services.

Having made situational analysis in details, highlighting points of weaknesses and strengths and defining actual needs, strategic plans were subsequently developed putting into practice the reforming infrastructure and human resources as well as partnerships between governmental, private and national sectors.

It gives me great pleasure to present this document. This system is in continuous reform, progressing incrementally, refining the knowledge base, and modifying concepts. This document is not the end product, but rather the first step of many others.

However, I hope it will help us towards our ultimate goal of a quality, effective, efficient, evidence based service to all Egyptians irrespective of geographical or social and economic barriers.

The document is a collaborative work of the Ministry of Health and Population staff, and the Sector for Technical Support and Projects on both central and peripheral levels. Work in this document is subjected to continuous assessment, operation research, many of the issues presented in this document will be updated in further version.

Table of Contents

Preface	Message from his Excellency	j
List of Figures	·	
List of Tables		
1- EYE PROBLEMS		
Trigeminal Neuralgia		
2- Dental Care & Oral Medicine 13	1- Eye Problems	
Trigeminal Neuralgia 13 Dental Care & Oral Medicine 12 Lip Swelling 12 Lip Pain 13 Gum Hypertrophy 14 Bleeding Gum 14 Purpura and Abnormal Bleeding 15 Colour of The Tongue 15 Pain in The Tongue 16 Papillae of The Tongue 16 Tongue Mass or Swelling 17 Furring of The Tongue 17 Taste Abnormalities 18 Stomatitis 18 Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3-Surgical Emergencies Burns and Scalds 35 Hemorrhage 36 Chocking 37	The Red Eye	9
Dental Care & Oral Medicine 13 Lip Swelling 13 Lip Pain 13 Gum Hypertrophy 14 Bleeding Gum 14 Purpura and Abnormal Bleeding 15 Colour of The Tongue 15 Pain in The Tongue 16 Papillae of The Tongue 16 Tongue Mass or Swelling 17 Furring of The Tongue 17 Taste Abnormalities 18 Stomatitis 18 Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies 26 Burns and Scalds 35 Hemorrhage 36 Chocking 37	2- DENTAL CARE & ORAL MEDICINE	
Dental Care & Oral Medicine 13 Lip Swelling 13 Lip Pain 13 Gum Hypertrophy 14 Bleeding Gum 14 Purpura and Abnormal Bleeding 15 Colour of The Tongue 15 Pain in The Tongue 16 Papillae of The Tongue 16 Tongue Mass or Swelling 17 Furring of The Tongue 17 Taste Abnormalities 18 Stomatitis 18 Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies 26 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Trigeminal Neuralgia	13
Lip Swelling 13 Lip Pain 13 Gum Hypertrophy 14 Bleeding Gum 14 Purpura and Abnormal Bleeding 15 Colour of The Tongue 15 Pain in The Tongue 16 Papillae of The Tongue 16 Tongue Mass or Swelling 17 Furring of The Tongue 17 Taste Abnormalities 18 Stomatitis 19 Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 22 Opisthotonus 24 3-Surgical Emergencies 24 3-Surgical Emergencies 35 Hemorrhage 36 Chocking 37	· ·	
Gum Hypertrophy 14 Bleeding Gum 14 Purpura and Abnormal Bleeding 15 Colour of The Tongue 15 Pain in The Tongue 16 Papillae of The Tongue 16 Tongue Mass or Swelling 17 Furring of The Tongue 17 Taste Abnormalities 18 Stomatitis 18 Mouth Pigmentation 19 Hyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies 26 Basic Life Support 28 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Lip Swelling	
Bleeding Gum 14 Purpura and Abnormal Bleeding 15 Colour of The Tongue 16 Papillae of The Tongue 16 Tongue Mass or Swelling 17 Furring of The Tongue 17 Taste Abnormalities 18 Stomatitis 19 Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies 26 Basic Life Support 28 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Lip Pain	
Bleeding Gum 14 Purpura and Abnormal Bleeding 15 Colour of The Tongue 16 Papillae of The Tongue 16 Tongue Mass or Swelling 17 Furring of The Tongue 17 Taste Abnormalities 18 Stomatitis 19 Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies 26 Basic Life Support 28 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Gum Hypertrophy	14
Colour of The Tongue 15 Pain in The Tongue 16 Papillae of The Tongue 16 Tongue Mass or Swelling 17 Furring of The Tongue 17 Taste Abnormalities 18 Stomatitis 19 Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies 25 Burns and Scalds 35 Hemorrhage 36 Chocking 37		
Colour of The Tongue 15 Pain in The Tongue 16 Papillae of The Tongue 16 Tongue Mass or Swelling 17 Furring of The Tongue 17 Taste Abnormalities 18 Stomatitis 19 Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies 25 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Purpura and Abnormal Bleeding	
Papillae of The Tongue 16 Tongue Mass or Swelling 17 Furring of The Tongue 17 Taste Abnormalities 18 Stomatitis 19 Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies 26 Burns and Scalds 35 Hemorrhage 36 Chocking 37		
Tongue Mass or Swelling 17 Furring of The Tongue 17 Taste Abnormalities 18 Stomatitis 19 Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies 26 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Pain in The Tongue	16
Furring of The Tongue 17 Taste Abnormalities 18 Stomatitis 19 Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies Basic Life Support 28 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Papillae of The Tongue	
Taste Abnormalities 18 Stomatitis 19 Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies 28 Burns and Scalds 35 Hemorrhage 36 Chocking 37	•	
Stomatitis 19 Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies 28 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Furring of The Tongue	17
Mouth Pigmentation 19 Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies Basic Life Support 28 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Taste Abnormalities	
Ptyalism 20 Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies Basic Life Support 28 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Stomatitis	
Myasthenia Gravis 20 Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies Basic Life Support 28 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Mouth Pigmentation	
Halitosis 21 Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies Basic Life Support 28 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Ptyalism	
Jaw Pain 22 Jaw Swelling 23 Opisthotonus 24 3- Surgical Emergencies 28 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Myasthenia Gravis	
Jaw Swelling	Halitosis	21
Opisthotonus 24 3- Surgical Emergencies 28 Basic Life Support 28 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Jaw Pain	
Opisthotonus 24 3- Surgical Emergencies 28 Basic Life Support 28 Burns and Scalds 35 Hemorrhage 36 Chocking 37	Jaw Swelling	
Basic Life Support		
Burns and Scalds 35 Hemorrhage 36 Chocking 37	3- Surgical Emergencies	
Burns and Scalds 35 Hemorrhage 36 Chocking 37	Racic Life Support	?ç
Hemorrhage		
Chocking		
G	<u> </u>	
THE STREET VICEOUS ARCHIVATOR	Intestinal Obstruction	

4- MEDICAL EMERGENCIES


Coma	44
Disturbed Level of Consciousness	
Syncope	46
Delirium	47
Epilepsy	
Precipitating Factors	
Convulsions	
Stridor	48
Acute Abdominal Pain	
5- Food Poisoning	
Food Poisoning	53
6- Minor Surgery	
Acute Abscess	57
Wounds	
Wounds for Urgent Referral	
Skin Lesions	
7- MANAGEMENT OF LIVER DISEASES	
Acute Liver Disease	61
Chronic Liver Disease	
Jaundice	
Prescribing in Hepatic Patients	
Role of The Liver in Dealing with Drugs	
Factors Affecting Bio-Transformation of Drugs	
Mechanisms of Impaired Drug Disposition in Liver Disease	
CYP ₄₅₀ Inducers	
Pharmacokinetic and Pharmacodynamic	
Effect of Liver Disease on Some Pharmocologic Agents	
Guideline Development Group Acknowledgements	71

List of Figures

No	Title	Page	
1	Flow-Chart Diagram for Symptoms of mouth disease		
2	Flow-Chart Diagram for D.D. of Lip Pain		
3	Flow-Chart Diagram for D.D. of Lip Swelling	13	
4	Flow-Chart Diagram for D.D. of Hypertrophy		
5	Flow-Chart Diagram for D.D. of Bleeding Gum	14	
6	Flow-Chart Diagram for D.D. of Pursuer and abnormal Bleeding	15	
7	Flow-Chart Diagram for D.D. of Papillae of The Tongue	16	
8	Flow-Chart Diagram for D.D. Tongue Mass or Swelling	17	
9	Flow-Chart Diagram for D.D. for Taste Abnormalities	18	
10	Flow-Chart Diagram for D.D. of Mouth Pigmentation	19	
11	Flow-Chart Diagram for D.D. of Ptyalism (Excessive Salivation)	20	
12	Flow-Chart Diagram for D.D. of Halitosis (Bad Odor)	21	
13	Flow-Chart Diagram for D.D. of Jaw pain	22	
14	Flow-Chart Diagram for D.D. of Jaw Swelling	23	
15	Flow-Chart Diagram for D.D. of Opisthotonus	24	
16	Diagram for D.D. of Position of the Victim	28	
17	Diagram for D.D. of Head tilt and chin lift	29	
18	Diagram for D.D. of Tongue obstructing airway; opening the airway by head tilt	29	
19	Diagram for D.D. of Mouth to mouth breathing	29	
20	Diagram for D.D. of Mouth to nose breathing	30	
21	Diagram for D.D. of Locating the carotid pulse	30	
22	Diagram for D.D. of Proper position of rescuer	31	
23	Diagram for D.D. of Finding proper hand position	31	
24	Diagram for D.D. of Hand position for external chest compressions	32	
25	Diagram for D.D. of Chest compressions	32	
26	Diagram for D.D. of Head tilt and chin lift for an infant	33	
27	Diagram for D.D. of In an infant check for the brachial pulse	33	
28	Diagram for D.D. of Heimlich maneuver for conscious	34	
29	Diagram for D.D. of Heimlich maneuver for unconscious	34	
30	Flow-Chart Diagram for D.D. of Coma or Stupor	45	
31	Flow-Chart Diagram for D.D. of Syncope	46	
32	Flow-Chart Diagram for D.D. of Delirium	46	
33	Flow-Chart Diagram for D.D. of Convulsions	47	
34	Flow-Chart Diagram for D.D. of Stridor	48	
35	Flow-Chart Diagram for D.D. of Acute Abdominal Pain	49	
36	Flow-Chart Diagram for D.D. of Jaundice	62	

List of Tables

No	Title	Page	
1	Variation between Infant, Child and Adult CPR		
2	Glasgow Coma Scale		
3	Bacterial Causes of Food Poisoning		
4	D.D of Jaundice	62	





Eye Problems

The Red Eye

Conjunctivitis: inflammation of the conjunctiva is the commonest eye problem seen in general practice:

- A. Infective Conjunctivitis (Bacterial or Viral)
- B. Gonorrheal in Ophthalmia Neonatorum
- C. Allergic Conjunctivitis

Treatment:

Treat with Topical Antibiotic Eye Drops and/ or Anti-allergic Eye Drops.

Chloramphenicol "Antibiotic Preparation":

Application:

- Eye drops (0.5%): Apply 1 drop at least every 2 hours and continue for 48 hours after healing.
- Eye ointment (1%): Apply either at night (if eye drops used during the day) or 3-4 times daily (if eye ointment used alone).
- Side-effects: Transient stinging

Corticosteroids or other anti-infiammatory preparations:

(e.g. Dexamethazone - Betamethasone eye drops and ointment)

Application: (For short term treatment of inflamation)

Frequent applications until condition is controlled then reduce frequency of application



- Refer all cases for ophthalmology opinion.

Signs of a Potentially Dangerous Red Eye:

- Decreased visual acuity
- Pain deep in the eye
- · Absent or sluggish pupil response
- Corneal damage, fluorescein staining, history of trauma.

Refer patient to be seen by specialist the same day if in doubt.

Other Causes of Red Eve

- Orbital cellulites
- Episcleritis

- Scleritis
- Subconjunctival hemorrhage
- Iritis
- Blephritis
- · Acute glaucoma
- · Foreign body
- Dry eye syndrome
- · Corneal ulceration
- · Corneal aberration
- Arc eye

Refer

Refer all Cases for Ophthalmology Opinion.

Saving Sight Changing Lives

Trachoma is one of the most common causes of blindness in the developing world, and is most commonly associated with areas of extreme poverty. It is caused by bacteria called Chlamydia Trachomatis which is very similar to the one that causes the sexually transmitted disease Chlamydia.

Infection ---> repeated conjunctivitis

The infection spreads in a variety of ways:

The discharge, the flies &via fingers and clothes.

Beyond the irritation caused by conjunctivitis, there is more serious consequence: each infection leads to a small amount of scaring to the cornea & conjunctiva. This scaring builds up over years of repeated infection until trichiasis sets in(eyelashes turn inwards) ----> repeated trauma to cornea ---> it becomes opaque ---> causing vision to decline & irreversible blindness.


How can Trachoma be treated?


Antibiotics

Facial cleanliness

Environmental change

Surgery for trichiasis








Figure "1": Flow-Chart Diagram for Symptoms of mouth disease




Figure "2": Flow-Chart Diagram for D.D. of Lip Pain

Trigeminal Neuralgia:

Paroxysmal attacks of severe unilateral lancinating pain along distribution of one of the branches of trigeminal nerve. Pain is precipitated by touching localized trigger zones. On examination, there are no motor or sensory changes.

Treatment: Vitamin B12 1000μ IM/d for 1 wk Carbamazepine 200mg tds.

If persistent refer to dentist

• Lip pain with ulceration & herpes zoster

→ see dermatology section

Refer undiagnosed cases




Figure "3": Flow-Chart Diagram for D.D. of Lip Swelling

Refer undiagnosed cases.




Figure "4": Flow-Chart Diagram for D.D. of Gum Hypertrophy




Figure "5": Flow-Chart Diagram for D.D. of Bleeding Gum

Bleeding Gum

- Bleeding Gums With No Splenomegaly or Rash:
 - o Good oral hygiene is a must
 - o Regular brushing
 - o Regular visit to dentist

• Bleeding Gums With Splenomegaly or

- o It indicates serious disease; you have to refer for proper evaluation and diagnosis.
- o Also see skin section.
- o If bleeding gum is associated with other bleeding ---> **Refer**





Figure "6": Flow-Chart Diagram for D.D. of Purpura and abnormal Bleeding

Refer

Purpura & abnormal bleeding

It indicates serious disease; you have to refer for proper evaluation and diagnosis.



- **In Iron deffciency anaemia:** confirmed by complete blood picture (CBC)
 - o ↓HB
 - o ↓ MCV
 - o ↓MCH
 - o ↓MCHC

Treatment:

1. Treat the underlying cause


- 2. Give oral iron to correct anemia, 1 tablet tds. & check CBC after 2 wks →↑ in HB 1g/wk. If no response refer
- 3. Replace iron stores: continue oral Iron 6 months after correction of anemia.

• In Vitamin B₁₂ Deficiency Anemia: Treatment:

Vit B12 1000µg IM twice for 3 wk, then every 3

months for the rest of the patients 'life.

- In case of central cyanosis, refer all cases for proper evaluation
- Leukoplakia is precancerous lesion, refer for excision
- Candida→ see dermatology section



• In Vitamin B₁₂ Deficiency Anemia:

Treatment:

Vit B_{12} 1000µg IM twice for 3 wk, then every 3 months for the rest of the patients 'life.

Prolonged antibiotics & menopause: give vit B complex

Referred pain: refer for further evaluation




Figure :7": Flow-Chart Diagram for D.D. of Papillae of The Tongue

• In Iron Deficiency anaemia: Confirm by Complete Blood Picture (CBC)

- o ↓HB
- o ↓MCV
- o ↓MCH
- o JMCHC

Treatment:

- 1. Treat the underlying cause
- 2. Give oral iron to correct anemia, 1 tablet t.d.s.
 & check CBC after 2 wks →↑ in HB 1 g/wk
 → If no response refer for parenteral iron therapy.
- 3.Replace iron stores: continue oral Iron 6 months after correction of anemia.


• In Vitamin B₁₂ Deficiency Anemia: Treatment:

Vit B₁₂ 1000µg IM twice for 3 wk, then every 3 months for the rest of the patients 'life.

Patchy desquamation: no clinical significance, and needs no treatment.

• Prominent Papillae (Scarlet Fever):

- o Oral penicillin 500mg four times daily for 10 days
- o Individuals allergic to penicillin can be treated by erythromycin 250mg four times for 10 days.



Scrotal tongue needs no treatment as it has no clinical significance.

Increased furring in Typhoid fever: ciprofloxacin twice daily for 14 days.




Figure "8": Flow-Chart Diagram for D.D. Tongue Mass or Swelling

Refer

Painful swelling must be referred to surgery for differential diagnosis Painless swelling referred for further evaluation & proper diagnosis.




Figure "9": Flow-Chart Diagram for D.D. Taste Abnormalities

Revise drug treatment of the patient to exclude drug cause Then examine mouth locally to exclude local cause, if present refer to dentist The remaining cases are referred for further evaluation & proper diagnosis




Figure "10": Flow-Chart Diagram for D.D. of Mouth Pigmentation

Fabry's Disease:

X-linked recessive disease leading to accumulation of glycophospholipids in lysosomes of liver, kidney, blood vesels and nervous system.

Refer for genetic counselling and identification of carriers.

In Addison's Disease:

Long term treatemnt with replacement glucocorticoids and mineralocorticoids.

Patient's advice:

o Carry a 'steroid card'

- o Wear a Medic-Alert bracelet
- o Keep an (up to date) ampoule of hydrocortisone at home


Peutz Jeghers:

Autosomal dominant disorder consists of mucocutaneous pigmentations and GI polyps.

Follow up every 2 years by x-ray & endoscopy

Acanthosis Nigricans:

Part of metabolic syndrome \rightarrow see diabetic section



Stomatitis

• Stomatitis: inflammation of buckal mucosa.

• Catarrhal in Flammation :

Red, dry, swollen buccal mucosa, covered with brownish secretion and foul smelling.

It occurs with poor oral hygiene, and with excessive smoking. Treatment: stop smoking

and proper mouth hygiene

• Aphthous :

Painful Ulcer

Treatment: local anesthetic

• Thrush:

White removable membranes see dermatology section

Vincent:

Fever, ulcers and foul odor broad spectrum antibiotic + flagyl

Then Refer All Cases.

Xerostomia

Mouth breather

Fear, anxiety

Fever

Dehydration

Drugs: atropin

Sjogren Syndrome:

keratoconjunctivitis sicca

- Idiopathic
- · Associated with collagen disease.
- Associated with lympho proliferative disorders.

Treatment: treatment of the cause

chewing gums, pilocarpin

Refer undiagnosed cases




Figure "11": Flow-Chart Diagram for D.D. of Ptyalism (Excessive Salivation)

Ptyalism (Excessive Salivation)

Revise drugs of the patient

Then examine mouth cavity, if abnormalities→ refer to dentist

Myasthenia Gravis:

Acquired condition is characterized by weakness and fatigability of proximal limb, ocular and bulbar muscles.

Refer for further evaluation

Cases of constant ptyalism refer for further evaluation

Pseudo Bulbar Palsy:

LMNL: Lower Motor Neuron Lesion of 9th., 10th., 11th., 12th., cranial nerves

Facial Palsy

Parkinsonism

Dementia

20

Halitosis (Bad Odor)




Figure "12": Flow-Chart Diagram for D.D. of Halitosis (Bad Odor)

Foul smelling → broad spectrum antibiotics and flagyl Refer undiagnosed cases

Bronchiectasis

Lung abscess




Figure "13": Flow-Chart Diagram for D.D. of Jaw pain

Jaw pain with abnormal mouth examination, refer to dentist

Trigeminal Neuralgia

Treatment: Vitamin B12 1000 μ IM / d for

1 wk

Carbamazepine 200mg t.ds.

If persistent refer to dentist

Cases with constant pain, refer for further evaluation

Herpes Zoster

Mumps




Figure "14": Flow-Chart Diagram for D.D. of Jaw Swelling

Paget's Disease
Osteitis Fibrosa Cystica
Acromegaly

Refer

Focal jaw swelling, refer to dentist Diffuse jaw swelling, refer to surgeon

Figure "15": Flow-Chart Diagram for D.D. of Opisthotonus

Refer

Acute Opisthotonus: Refer to Emergency Unit

Chronic Recurrent Opisthotonus:

Epilepsy: In case of epilepsy:

- Emergency Measures:
 - o maintain airway patent
 - o Avoid tongue biting
 - o Exclude hypoglycemia
 - o If prolonged seizure >3 min give IV diazepam
- Avoid Precipitating Factors:
 - o Pyrexia: in children <5 yrs.

- o Sleep deprivation
- o Photosensitivity: flashing lights or flickering
- o Drugs: tricyclic antidepressants, phenothiazines
- o Withdrawal of anticonvulsant drugs
- o Metabolic abnormalities

Refer for EEG & CT


In case of hysterical seizures: Reassure and refer to a psychiatrist

In the other cases refer for proper diagnosis.

Tetanus

Meningitis

24





Emergency Guidelines for Family Physicians

Topics:

- · Basic life support
- · Burns & scalds
- Shock
- Hemorrhage
- Fractures
- · Domestic accidents
- · Acute abdomen
- Intestinal obstruction
- Foreign body airway obstruction (choking)

Basic Life Support:

Deffnition:

It is the phase of emergency cardiac care that either prevents circulatory or respiratory arrest by prompt recognition and intervention, or externally support with cardiac pulmonary resuscitation (CPR) the circulation & respiration of a victim of cardiac and /or respiratory arrest.

The major objective of CPR is to provide oxygen to brain and other vital organs until appropriate, definitive medical treatment can restore normal heart & ventilation.

The victim whose heart & breathing have stopped for less than 4 minutes has an excellent chance for full recovery if CPR is administered immediately, after 4-6 minutes without circulation, brain damage may occur.

Sequence of Basic Life Support

1. The Assessment Phase (ABC Assessment)

- A. Airway: to determine unresponsiveness.
- B. Breathing: to determine breathlessness.
- C. Circulation: to determine pulselessness.

Assessment also involves observing and interacting with the victim. If unresponsiveness is established, activate the emergency medical system immediately.

2. Activate Emergency Medical System by Calling The Local Emergency Telephone Number.

3. Cardiopulmonary Resuscitation:

CPR includes the basic rescue skills, the victim must be supine and on a firm flat surface to help increase the blood flow to the brain. If the victim is lying face down, the rescuer must roll the victim as a unit "log roll" so that the head, shoulders & torso move simultaneously without twisting.



Figure "16": Position of the Victim

A. Airway

The first action for successful resuscitation is immediate opening of the airway, the tongue and epiglottis

are the most common cause of airway obstruction in the unconscious victim . Since the tongue is attached to the lower jaw , tilting the head back and moving the lower jaw forward to open the airway (head tilt / chin lift maneuver , figures 17&18)




Figure "17": Head tilt and chin lift

Basic Life Support




Figure "18": Tongue obstructing airway; opening the airway by head tilt

B. Breathing

When breathing stops , cardiac arrest and death quickly follow . Breathing is assessed by placing your ear over the victim's mouth and nose while maintaining an open airway .

While observing the victim's chest:

Look ... for the chest to rise and fall

Listen ... for the air escape during inhalation

Feel ... for the flow of air

Mouth to mouth breathing is the quickest way to get oxygen into the victim's lung Or mouth to nose breathing, this technique is more effective in some cases than mouth to mouth technique especially when

28

the mouth can not be opened (trismus) or seriously injured (figures 19 & 20)



Figure "19": Mouth to Mouth Breathing

Basic Life Support



Figure "20": Mouth to nose breathing

C. Circulation

Assessment of circulation to determine pulselessness.

Cardiac arrest is recognized by pulselessness in the large arteries of the unconscious victim (Carotid artery - figure 6).

This assessment should not take more than 5-10 seconds.




Figure "21": Locating the carotid pulse




Figure "22": proper position of rescuer




Figure "23": Finding proper hand position



Figure "24": Hand position for external chest compressions




Figure "25": Chest compressions

Two - Rescuer CPR:

One rescuer moves to head, opens the airway and checks the pulse, while the other rescuer locates the area of chest compression and finds the proper hand position, this should take 5 seconds only. If there is no pulse, the ventilator person gives 2 breaths and compression counting 5 times (2 Breaths-5 Compressions).

CPR in Infants and Children:

Cardiopulmonary resuscitation and air way obstruction management for children over 8 years old is the same for the adult. but for small children (from 1-8 years) and

infants (Less than 1 year) there are important differences in techniques as a result of small size and physical immaturity of the age group. These differ in giving breathing (1.0-1.5sec/infantandchild). Chestcompression depth (1.3 to 2.5 cm in infants and 2.5 to 3.8 cm in child). Chest compression in infants is performed with 2 fingers on the lower third of the sternum and in a child with the heal of one hand on the lower third of the sternum.

Figure "26": head tilt and chin lift for an infant




Figure "27": In an infant check for the brachial pulse

Basic Life Support Respiratory Emergency:

Definition: Respiratory Emergency is that condition where normal breathing stops OR oxygen intake markedly falls that it becomes insufficient to support life.

Causes: Upper air way obstruction can cause unconsciousness and cardiopulmonary arrest An unconscious victim can develop air way obstruction when the tongue falls backward into the pharynx, regurgitation of stomach contents into the pharynx, bleeding from head and facial injuries may also obstruct the upper airway, particularly if the victim is unconscious. Foreign body airway obstruction usually occurs during eating.

Management of Obstructed Airway:

The Heimlich maneuver (sub diaphragmatic abdominalthrusts) (Figures 28,29) is recommended for relieving foreign body airway obstruction. By elevating the diaphragm the Heimlich maneuver can force air from the lungs to create an artificial cough intend to expel a foreign body obstructing the airway. Each individual thrust should be administered with the intent of relieving the obstruction. It may be necessary to repeat the thrust multiple times during each sequence to clear the airway.



Figure "28": Heimlich maneuver for conscious



Figure 29: Heimlich maneuver for unconscious

Summary

Cardiopulmonary Resuscitation (CPR) is a holding action for sudden cardiac or respiratory arrest until more advanced life support care can be made available.

CPR involves a combination of mouth to mouth rescue breathing and chest compression. It keeps some oxygenated blood flowing to the brain and other vital organs until appropriate medical treatment can restore normal heart action. The major factor that influences success is Speed. The highest success rate has been achieved whn PR was started within 4 minutes or less. In order to ensure fast action it is important to recognize the need for immediate action and perform the steps of CPR quickly and efficiently.

Cardiopulmonary Resuscitation includes three basic rescue skills, the ABCs of CPR: Airway, Breathing and Circulation to Assess the Victim's condition and to support life. This table Summaries the variation between Infant, Child, and Adult as regard CPR:

Table. 1: Variation between Infant, Child, and Adult CPR

Item	Infant 0-1 Year	Child 1-8 Years	Adult>8 Years
Shake and Shout	Yes	Yes	Yes
Call for Help	Yes	Yes	Yes
Position Victim	Yes	Yes	Yes
Open Airway	Yes	Yes	Yes
Look, Listen and Feel for	Yes	Yes	Yes
breath			
Two breaths	Yes	Yes	Yes
Check Pulse	Brachial	Carotid	Carotid
Activate EMS	Yes	Yes	Yes
Locate Hand Position	Lower third sternum	Lower third	Lower third
		sternum	sternum
Compress with	2-3 fingers	Heal of one Hand	Heal of one Hand
_	_		other Hand on Top
Compression Depth	1.3-2.5 cm	2.5-3.8 cm	3.8-5 cm
Compression / minute	At least 100	80-100	80-100
Compression /	5-1	5-1	5-2 or 5-1
Ventilation Ratio			
Breathing	2 breaths 1-1.5	2 breaths 1-1.5	2 breaths 1.5 -2
	seconds each.	seconds each.	seconds each.
	20/minute	20/minute	10- 12/minute

Burns and Scalds

Burn types:

- · Scald/ steam
- Contact
- Chemical/caustic
- Electrical
- Radiation

Clinical Assessment:

- Cause, size, and thickness.
- Use rule of nines.
- Partial thickness burns are painful, red & blistered.[Do not remove or rupture blisters].
- Full thickness burns are painless and white or gray.

Rule of Nines:

palm	1%
arm(all over)	9%
leg (all over)	18% (14%children)
front	18%
back	18%
head(all over) genitals	9% (14% children) 1%

Action

- Ignore areas of erythema only.
- Remove clothes from affected area.
- Place under cold running water for 10 minutes or until pain is relieved.
- Do not burst blisters.
- · Give analgesia.
- Check tetanus immunity give immunization and/or prophylaxis as necessary.
- Apply silver sulfadiazine or sterile Vaseline -

impregnated gauze and non adherent dressing and follow up, every 1-2 days (for healing & infection).

Refer

Refer The Following Cases:

- 1. Cases of more than 10% -children or 15% adults.
- 2. Burns affecting: face (eyes & ears), neck, both hands, both feet & perineum.
- 3. Type of burn: electrical, chemical & smoke inhalation.
- 4. Presence of associated injuries: soft tissue trauma, fractures & head injury.
- 5. Presence of medical problems: diabetes, heart disease, pulmonary disease & ulcer history.
- 6. Social problems such as suspected child abuse or neglect, self inflicted burns & psychological problems.

Prevention of Scalds & Burns

Health education is very important (in occupational & domestic accidents)

Domestic Accidents

- The leading causes of accidental deaths & injuries in the home for all age groups are:
- Falls
- Fires
- Burns
- · Poisoning
- Suffocations
- All age groups are affected but children &


elderly are more susceptible (Risk group).

- Risk factors for accidents in the elderly group:
 - · Chronic illness
 - · Neurological abnormalities
 - · Psychological disturbances
 - · Poly pharmacy
 - · Visual impairment
 - Osteoporosis
 - · Gait abnormalities

Domestic Accidents may present as:

Wounds, burns, scalds, fractures, eye injuries, shock, hemorrhage etc...

Action :-



Shock

Definition: the presence of inadequate tissue perfusion & oxygenation.

Compensated shock	Uncompensated shock
Vital organ perfusion is maintained by compensatory mechanisms	Associated with impairment of tissue perfusion when compensatory mechanisms fail

Types of Shock:

- · Hypovolemic shock
- · Cardiogenic shock
- Tension Pneumothorax
- Neurogenic shock
- Anaphylactic shock
- Septic shock
- Endocrinal shock

Recognizing Shock Clinical findings Vital sings Peripheral perfusion Color Mental status No reliable laboratory tests

Hypovolemic shock:

Caused by diminished blood volume secondary to loss of:

• blood, as in external or internal hemorrhage

(hemorrhagic shock)

- plasma, as in burns and peritonitis
- fluids, as in severe vomiting, diarrhea, intestinal obstruction or in a high output intestinal fistula.

Recognizing Shock Hemorrhagic Shock

Most common in trauma

Not common in isolated head injury

Aggressive management improves outcome

Recognizing Shock Cardiogenic shock

Myocardial dysfunction Cardiac tamponade

Recognizing Shock Tension pneumothorax

Clinical diagnosis

Emergent needle decompression

Recognizing Shock

Neurogenic Shock

Loss of sympathetic tone

Hypotension without tachycardia No cutaneous vasoconstriction

Recognizing Shock

Septic Shock

Uncommon in acute trauma

Warm skin

Wide pulse pressure

Management:

Shock Management

Airway

Breathing

Circulation Disability

Exposure

Shock Management

Circulation

Control external bleeding

Obtain vascular access

Fluid resuscitation(isotonic electrolyte solution-

Ringer's lactate-Normal saline

Adult 1-2 liters

Children 20 ml/kg

Do ABCs and transfer into emergency unit in nearest hospital.

Hemorrhage

Definition: It is the escape of blood from a vessel damaged by injury or disease.

Types of Hemorrhage:

- 1. Traumatic hemorrhage
- · primary occurs immediately after trauma
- · secondary is complication of sepsis
- · reactionary occurs within the first 24 hrs, after cessation of the primary
- 2. Pathological hemorrhage
- inflammatory, neoplastic, etc.
- 3. Spontaneous hemorrhage
- · defect in haemostatic mechanisms

Clinical types

Internal bleeding (concealed) e.g.

- intracranial
- · haemothorax
- haemoperitoneum
- · haempericardium
- · retroperitoneal

External bleeding (revealed) e.g.

- wound • ulcer
- fungating tumor
- from natural orifice: o Epistaxis
- Haemoptysis
- · Haematemesis
- Haematuria
- o Melaena • Metrorrhagia

Management of hemorrhage:

Among all surgical emergencies, hemorrhage constitutes the most urgent problem as it rapidly threatens the patients' life & predisposes to serious local & general complications.

The principles of management of all forms of hemorrhage consist simply of:

- Immediate control of bleeding hemorrhage)
- Timely replacement of the lost blood (restore blood volume)

In practice, the management of bleeding from external wound consists of 3 phases:

First-aid treatment: 3 P's

(Packing, Pressure & Position)

Digital compression Pressure bandage or by sphygmomanometer Elevation of the limb above the level of the heart

- 1. Resuscitation by anti- shock measures
- 2. Definitive control of bleeding by operation.

Chocking

1. Choking-Adult

If blockage of airway is only partial, the victim can usually be able to dislodge the foreign body by coughing, but if obstruction is complete urgent intervention is required to prevent asphyxia.

Conscious Adult:

Carry Out Back Blows:

- Stand beside & slightly behind the victim
- Support the chest with one hand & lean the victim forewords
- Give up 5 sharp blows between the scapulae with the heel of the other hand; each blow should be aimed to relieve the obstruction
- · If the back blows fail, carry out abdominal thrusts.

Give 5 Abdominal Thrust (Heimlich Maneuver):

- · Stand behind the victim
- Wrap your arms around victims waist
- Make a fist with one hand & place the thumb side just above victim's navel & below the tip of the sternum (xiphoid process)
- · Grasp fist with your other hand
- Press fist into victim's abdomen with quick upward thrusts
- Each thrust should be a separate& distinct effort to dislodge the object
- · After every 5 abdominal thrusts check the victim's mouth, coughs up object, or start to breath or cough forcefully.



Unconscious Adult:

A -Give up to 5 abdominal thrusts with the victim supine on the floor:

- Straddle victim's thighs
- Put heel of one hand against middle of victim's abdomen slightly above navel & well below sternum notch (fingers of hand should point towards victim's head)
- Put other hand directly on top of the first hand
- Press inward & upward using both hands with up 5 quick abdominal thrusts
- Each thrust should be distinct & a real attempt made to relieve the airway obstruction.
- Keep the heel of hand in contact with abdomen between thrusts.

B-Perform finger sweep:

- Use your thumb & fingers to grasp victim's jaw & tongue, lift the tongue away from back of throat & away from the foreign body
- Or use cross finger method by crossing the index finger & thumb and pushing the teeth apart.
- With index finger of other hand, slide finger down along the inside of one cheek deeply into mouth & use a hooking action across the

other cheek to dislodge foreign object

• If foreign body comes within reach, grab & remove it, do not force object deeper

C- Sequence the cycle:

- Give 2 rescue breaths
- Do up to 5 abdominal thrusts
- Finger sweep

Choking-Child

(1-8 years)

- Look into mouth
- Remove foreign body only if seen with finger sweep
- Give up 5 abdominal thrusts (Heimlich maneuver)

Choking Infant

(Under 1 year)



Note

A. If Infant is Conscious: (Can Cough, Breathe or Cry)

- 1. Shout for help
- 2. Give up to 5 back blows:
 - Lay infant face down over your forearm with head lower than his/her chest
 - Brace your forearm & infant against your thigh
 - Give up to 5 distinct & separate back blows between shoulder blades by the heel of your hand
- 3. Give up to 5 chest thrusts

By supporting the back of infant's head & start CPR

- 4. Repeat:
 - up to 5 back blows
 - up to 5 chest thrusts

Note

B. If Infant is Unconscious:

1. Give 2 slow breaths

- Open the airway with headtilt/ chin lift method
- Seal your mouth over the infants mouth & nose
- Give 2 slow mouth to mouth breaths 1-1½ seconds each
- 2. Activate emergency medical system for help
- 3. Give up to 5 back blows
- 4. Give up to 5 chest thrusts
- 5. Check mouth for foreign body (if object is seen use finger sweep)
- 6. Repeat the cycle until help arrives or object is expelled.

Fractures & Extremity Trauma

Identify

- Life threatening injuries
- Limb threatening injuries
- · Disabling injuries
- · Less serious injuries

38

How life threatening?

- · Associated injuries
- · Hemorrhage
- · Crush syndrome
- · Overwhelming infection
- Multiple injuries could add to risk of multi- system organ

How limb threatening?

- · Vascular injury
- · Compartmental syndrome
- · Complete disorganization

Attention

- · Documentation is important
- Do not allow the limb injury distract you, There may be a more serious & important injury
- · Remember occult injuries
- Set priorities in assessment & management
- · Activate Emergency medical system

Primary survey & Resuscitation

- Airway
- · Breathing
- Circulation
- Disability
- Exposure
- Splint
- · Restore distortion
- · Wound care
- Refer to emergency unit of the nearest hospital

NB: X ray is not always urgent

Intestinal Obstruction

Intestinal obstructions is a common surgical emergency, because of its serious nature, it demands early diagnosis & speedy relief

Types:


A Dynamic Obstruction Peristalsis ceases & no.

propulsive waves occurs

Dynamic Obstruction

Peristalsis is working against obstructing agent (in the lumen, the wall or outside the wall)


Clinical Types:



- · Acute intestinal obstruction favors small gut, with immediate central severe, colicky abdominal pain, early vomiting, central distension, and constipation.
- Chronic obstruction favors the large bowel, with lower abdominal colic at first & absolute constipation. Distension comes later, and involves the periphery.
- Acute on top of chronic obstruction; spreads from the large bowel to involve the small intestine.It gives rise to pain & constipation on a variable time scale, later followed by general distension & vomiting.



· Refer all cases of intestinal obstruction to nearest hospital.





Management of Medical Emergencies

Unconsciousness and Coma

- Glasgow coma scale
- Disturbed level of consciousness
- Delirium
- Syncope
- Convulsions
- Stridor
- Abdominal pain: Acute

Unconsciousness and Coma

- Consciousness means a state of wakefulness with awareness of self and surroundings.
- Clouding of consciousness means reduced wakefulness and/or self-awareness.
- Confusion is the state of altered consciousness

- in which the subject is bewildered and misinterprets his or her surroundings.
- Sleep is the state of normal mental and physical inactivity from which the subject can be aroused.
- Stupor is an abnormal, sleepy state from which the subject can be aroused by stimuli, applied vigorously or repeatedly.
- Delirium is a state of high arousal in which there is confusion and often visual hallucination.
- Coma is a state of unarousable unresponsiveness (The Glasgow coma Scale for grading coma).
- Syncope (drop attack) is a transient loss of consciousness (fall to ground) for less than 2 minutes and recovered when patient lies down.
- Glasgow Coma Scale

Table. 2: Glasgow Coma Scale

Eye opening (E)	Score	Verbal response (V)	Score
Spontaneous	4	Orientated	5
To speech	3	Confused conversation	4
To pain	2	Inappropriate words	3
No response	1	Incomprehensible sounds	2
		No response	1
Motor response (M)			
Obeys	6		
Localizes	5		
Withdraws	4		
Flexion	3		
Extension	2		
No response	1		
Class Constitution France			

Glasgow Coma Scale = E + M + V (GCS minimum = 3 : maximum = 15)




Figure "30": Flow-Chart Diagram for D.D. of Coma or Stupor

Disturbed Level of Consciousness

- First you have to exclude hypoglycemia.
- If you suspect hypoglycemia give oral or IV glucose

Coma:

is a state of unarousable unresponsiveness (The Glasgow coma Scale for grading coma).

Stupor:

is an abnormal, sleepy state from which the subject can be aroused by stimuli, applied

vigorously or repeatedly.

Note

After exclusion of hypoglycemia, in all patients examine & assess Glasgow Coma Scale

- Ensure patent airway
- Put IV line with 5% glucose to keep veins open
- Refer to emergency unit




Figure "31": Flow-Chart Diagram for D.D. of Syncope

Syncope:

(drop attack) is a transient loss of consciousness (fall to ground) for less than 2 minutes and recovered when patient lies down.

Immediate management:

lay patient down & lift legs, Patient will recover consciousness

Then refer for further evaluation




Figure "32": Flow-Chart Diagram for D.D. of Delirium

Delirium

Delirium is a state of high arousal in which there is confusion and often visual hallucination.

After exclusion of hypoglycemia, in all patients examine & assess Glasgow Coma Scale

Ensure patent airway

Put IV line with 5% glucose to keep veins open Refer to emergency unit




Figure "33": Flow-Chart Diagram for D.D. of Convulsions

In Case of Epilepsy:

Note

Emergency Measures:

- Maintain airway patent
- Avoid tongue biting
- Exclude hypoglycemia
- If prolonged seizure >3 min give IV diazepam

Avoid Precipitating Factors:

• Pyrexia: in children <5 yrs.

- Sleep deprivation
- Photosensitivity: flashing lights or flickering
- Drugs: Tricyclic Antidepressants, Phenothiazines
- Withdrawal of anticonvulsant drugs
- Metabolic abnormalities

Refer for EEG & CT

In Case of Hysterical Seizures:

Reassure and refer to a psychiatrist

In the other Cases Refer for Proper Diagnosis.

46




Figure "34": Flow-Chart Diagram for D.D. of Stridor

Stridor: is difficulty in respiration (inspiratory) or due to pathology in upper respiratory tract.




Figure "35": Flow-Chart Diagram for D.D. of Acute Abdominal Pain

Acute Abdominal Pain

In medical causes of acute abdomen there is NO rigidity or rebound tenderness.

- **N.B.** Familial Mediterranean Fever is characterized by recurrent attacks of
- Fever
- Arthritis: monoarticular

48

- Serositis: abd.pain due to peritonitis or pleurisy
- Attacks last for up to 1 week

Refer to confirm diagnosis

Appendicitis produces more gradual onset of pain and pain may be made worse by movement.

Vomiting may accompany any acute abdominal pain but, if persistent, it suggests an obstructive lesion of the gut.

All other cases with rigidity and /or rebound tenderness should be referred to surgical emergency unit.

Other rare medical Cause of Acute Abdomen are acute propheria and familial hypertriglyceridemia.

Refer

A sudden onset of severe pain suggests:

- Perforation e.g.doudenal ulcer
- Rupture e.g. of an aneurysm
- Torsion e.g. of an ovarian cyst
- Acute pancreatitis

Refer immediately to emergency unit

Colicky pain can be due to an obstruction of Gut

- Biliary system
- · Urogenital system
- Or uterus.

These will probably initially require conservative management along with analgesics and antispasmodic.

If colicky pain changed into constant pain \rightarrow inflammation supervene. This will be supported by:

- Raised temperature.
- Tachycardia
- And/or raised white cell count

Add broad spectrum antibiotics, IV line and transfer to emergency unit

Back Pain Suggests:

- · Pancreatitis
- · Rupture of an aortic aneurysm

- Urinary tract disease
- **Diabetic Ketoacidosis** (refer to Diabetic section)
- Myocardial infarction: refer to chest pain & IHD section
 - o Give sublingual nitrate tablet every 5 min for 3 tablets
 - o Chew aspirin tablet
 - o Refer to emergency unit

• Sickle Cell Crisis:

- o IV fluid
- o Oxygen
- o Antibiotics
- o Adequate analgesia
- o After attack give pneumococcal vaccine
- o Hemophilus influenza vaccine
- o Refer to further evaluation

Scorpion Bite:

- 1.analgesic
- 2.antivenom

Snakes Bite:

- 1. observation for 12-24h
- 2.no excision of bite area
- 3.no arterial tourniquet
- 4. diazepam for anxiety
- 5.I.V fluids & volume expander for hypotension
- 6. Antivenoms as early as possible by slow IV
- 7. usually there is allergy give adrenalin (1 in 1000) & continue antivenoms
- 8. antibiotic if wound is infected
- 9. antitetanous prophylaxis

Tetanus:

- 1.clean wound & debridment
- 2.give tetanus Ig 250u + tetanus toxoid "booster"
- 3.if not vaccinated before: give tetanus toxoid:

 day zero
 0.5ml IM

 8 wks
 0.5ml IM

 6-12 m
 0.5ml IM

booster every 5 years





Food Poisoning

Disease of an infective or toxic nature caused by consumption of food and water Food poisoning may also be caused by organic & non-organic toxins.

Table 3: Bacterial Causes of Food Poisoning:

Organism	Source	IP	Symptoms	Diagnosis	Recovery
Staph.aureus	Contaminated food & water	2-4 h	Diarrhea, vomiting & dehydration	Culture organism in vomitus of food	< 24 h
E.coli	Cattle meat, milk	12-48 h	Watery diarrhea (± bloody diarrhea)	Stool culture	10-12 days
Bacillus cereus	Contaminated food	1-6 h	Diarrhea, vomiting & dehydration	Culture organism in feaces and food	Rapid
Clostridium perfringes	Contaminated food	8-22 h	Watery diarrhea & cramping pain	Culture organism in faeces and food	2-3 days
Clostridium botulinum	Bottled or canned food	18-24 h	Neuromuscular paralysis	Demonstrate toxin in food or faeces	10-14 days
Salmonella	Cattle and poultry eggs, meat	12-48 h	Diarrhea, vomiting & fever	Stool culture	3-6 days, up to 2 weeks
Campylobacter	Cattle and poultry meat, milk	48-96 h	Diarrhea ± blood, fever, malaise & abdominal pain	Stool culture	3-5 days
Shigella	Man- contaminated food& water	24-48 h	Watery, bloody diarrhea	Stool culture	7-10 days


Prevention:

Improvement in food handling & preparation

Treatment:

Give oral fluid + empirical antibiotic

If patient is dehydrated IV line \rightarrow refer to secondary care





Minor Surgery

Guidelines of Minor Surgery for Family physician

- 1. Assess the case (history taking, physical examination...)
- 2. Follow basic principles of surgery:
 - Asepsis
 - Haemostasis
 - Local anesthesia
 - Suturing
 - Cautery
- 3. Follow up for healing & infection
- 4. Referral to Hospital for other surgery requiring general anesthesia & admission over night.

Types of Minor Surgery:

- 1. Nail extraction
- 2. Abscess drainage
- 3. Cut wound suturing
- 4. Male circumcision
- 5.Removal of skin lesion (sebaceous cyst, dermoid cyst, callosity, warts & small subcutaneous lipoma)

Acute Abscess Drainage:

- Once pus forms, the only treatment should be free drainage.
- Incision & drainage is the standard method..
 The incision should be situated over the pointing part of the abscess & should be dependent; otherwise a counter incision over a dependant area is necessary.
- The loculi inside the abscess are broken to ensure adequate free drainage.
- A specimen of pus could be sent for culture & sensitivity.
- A rubber drain or a wick of gauze is left protruding from the abscess cavity. **The drain serves two purposes**; it allows escape of discharge, and prevents premature closure of the skin wound before abscess cavity is obliterated by granulation tissue.
- Adequate free drainage is usually enough for most abscesses and antibiotics are required in only in overwhelming infections or in immunocompromised patients.
- In dangerous area, e.g. the neck, a sinus forceps is used to open the abscess after skin incision (Hilton's method) this avoids injury of important structures in the area.

Wounds

1. Assess the wound (history, physical examination).

History: ask about type of trauma, onset, duration, and treatments tried etc..

Physical Examination

- Examine while using sterile technique
- Assess the size, the depth of the wound
- Explore for foreign body
- Assess the circulation, sensation, and movement distal to the wound
- Palpate the underlying bones & parts surrounding the wound site.
- Assess the range of motion & strength against resistance of all body parts surrounding the wound.

2. Preparation of wound site

- Clip hair, do not shave surrounding hair as close to skin surface as possible. Eye brow should never be shaved.
- 3. Wound cleaning with normal saline
- 4. Prevention of tetanus is important (If person is immunized ---> give booster dose of the toxoid, if not immunized give immunoglobulin & start immunization)
- 5. After preparing the wound, decide whether to apply sutures

Refer

Wounds for Urgent Referral:

- Avulsion flaps: results in a sizable amount of non viable tissues that may not be apparent initially
- Amputated fingers if properly reserved can be re-implanted
- Stab wounds of the neck and the abdomen, even if apparently simple very commonly hide a serious injury
- Wounds involving nerve, tendon, or bone damage
- Facial and hand wounds if not small and /or superficial

Special Wound Conditions:

- · Old wounds
- Non accidental violent injury e.g. abused children, mentally retarded negligence, domestic accidents against women & self inflicted wounds.

In wounds Antibiotics

Prophylactic antibiotics should be given in the

- · Most mammal bites
- · Puncture wounds
- Patients with cardiac valve disease or implants
- Heavily contaminated wounds
- Wounds with delayed treatment
- Wounds with tissue necrosis

Removal of Skin Lesions

- Ensure that you have practiced the techniques
- Better remove benign lesions- refer suspicious lesions to a specialist
- Take care with lesions on the face & lip, the scare may be very noticeable
- Explain to the patient the possible complications & possibility of keloid
- Clean & anaesthetize the area
- Make incision in the skin lines if possible
- Send all excised lesions for histopathology, place in formalin & carefully label the site and side.
- Be careful with diabetic patients.

Sebaceous (Epidermoid) Cyst

Complete excision of the cyst with an ellipse of overlying skin containing the punctum is performed to avoid recurrence.

Dermoid Cyst

- The only way of treatment is by surgical excision.
- In children with a dermoid cyst in the scalp it is better to wait until closure of the skull sutures because some cysts may have communication with dura.
- Surgery for a dermoid cyst is more difficult than that for a sebaceous cyst, because a dermoid cyst is deeper.

Callosity

The patient should avoid the causative agent if possible. The callosity should be shaved with a razor blade & repeatedly painted with an ointment containing salicylic acid.

Corn

A corn is similar to a callosity but a combination of friction & pressure. Treatment is the same. excision & sutures are rarely necessary.

Wart

• This is due to virus infection.

Treatment Options Are:

- Curettage & Diathermy
- Repeated application of glacial acetic acid

Lipoma

A subcutaneous lipoma is usually excised for cosmetic reasons. The operation is easy & depend on enucleation of the tumor from the inside its capsule.

Circumcision

Circumcision is the most commonly performed operation for Egyptian males

There in nothing called female circumcision , it is forbidden & considered female genital organ mutilation.

Indications:

- Circumcision can be done at any age, better after 40 days post delivery
- It can be performed without anesthesia in the first two months
- Local infiltration an esthesia at age 2-6 months (fine needle & injecting 3-4 ml xylocaine 2% without adrenaline in the midline spot at the base of the shaft of the penis to block the dorsal nerve of the penis)
- Age 6 months & older give general anesthesia

Contraindication:

- · Bleeding disorders
- Hypospadius
- Epispadius

Postponement:

- · Presence of inguinal hernia
- Premature & unstable child
- Presence of local infection or extensive rash (napkin rash)

Techniques


- Aseptic preparation
- The method depends on the age of the patient:
- Crush- excision is the method of choice in infants & children
- Dissection- excision is necessary in adolescents & adults

Do not forget

Do not forget parent / patient education

Think in complications (post-operative bleeding, stricture injury of the glans penis & external urethra and gangrene of the glans) Try to avoid them.

58





Management of Liver Diseases

Liver Disease

- Acute liver disease
- Chronic liver disease
- Jaundice

Acute Liver Disease

Symptoms of Acute Liver Disease:

The patient may be:

Asymptomatic and anicteric or

Symptomatic: generalized symptoms of malaise, anorexia, fever then jaundice appear.

Signs of Acute Liver Disease:

Jaundice

Enlarged tender liver

Pale stool

Dark urine

Treatment:

If Hepatitis A:

No specific treatment.

Bed rest till liver enzymes return normal.

Small frequent light diet.

If Hepatitis B:

Prevention & prophylaxis:

- Standard safety precautions
- · Avoid risk factors
- Vaccination: see vaccination section
- Active immunization:
 - o Combined prophylaxis (vaccination &immunoglobulin)
 - o Staff with accidental needle-stick injury
 - o All newborn babies of HBs-positive patients.

Treatment as in Hepatitis A

If Hepatitis C:

- Prevention & Prophylaxis:
 - o Standard safety precautions.
 - o Avoid risk factors

Treatment as in Hepatitis A

In case of acute liver disease, refer patient for liver functions & viral markers

Chronic Liver Disease

Symptoms of Chronic Liver Disease:

The patient may be:

Asymptomatic, or

Nonspecific symptoms particularly fatigue, or Specific symptoms:

- Right hypochondrial pain
- Abdominal distension due to ascites
- Ankle swelling due to fluid retention
- · Hematemsis and melena
- Pruritis due to cholestasis
- Gynecomastia, loss of libido and amenorrhea
- Confusion and drowsiness due to neuropsychiatric complications (hepatic encephalopathy)

Signs of Chronic Liver Disease:

- May be normal, or
- Spider naevi over chest, upper body (above nipple line)
- · Palmer erythema
- Clubbing
- Duptuytren's contracture
- · Parotid enlargement
- · Shrunken liver
- Splenomegaly
- Gynecomastia, testicular atrophy
- Hepatic flap
- Foetor hepaticus
- Drowsy and coma
- Ascites, dilated veins on abdomen, lower limb oedema

In case of chronic liver disease, refer patient for diagnosis by for liver functions test & abdominal sonar.

Treatment:

- Salt restriction
- Avoid aspirin & NSAIDS

No treatment will arrest or reverse cirrhotic changes.

In case of complications, refer.

Jaundice

Definition yellowish discolorations of skin and mucous membrane due to \(\gamma \) serum bilirubin

	Hemolytic	Hepatocellular	Obstructive
Causes	Hemolytic anaemia	Acute: Viral hepatitis Toxic: gold CCL4 Chronic: Ch. Hepatitis Cirrhosis	Extra hepatic : Stone Malignancy Hepatic: Drugs:anabolic steroids, largactil, Cholangitis
Depth	Mild (lemon)	Moderate (orange)	Deep (olive green)
Bilirubin	↑ Indirect	↑ Both	↑ Direct > 20%
Urine	- Normal on voiding - Darken on standing	- Slightly dark	- Dark, frothy
Stool	Dark brown	Slightly pale	Clay colour
Itching	-	+	+++
Abd.Pain	- In acute hemolytic crisis - In scickle painful crisis	- In acute hepatitis	- Biliary stones - Malignancy
Associated	- Anaemia - Splenomegaly - + Leg ulcers	- Chronic liver cell failure - Portal hypertension	- Brady cardia - Coagulation defects - Fat malabsorption
Liver function tests	Normal	Impaired	- Early: normal except \\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \

Jaundice




Figure "36": Flow-Chart Diagram for D.D. of Jaundice

Prescribing in Hepatic Patients

- The main site of biotransformation is the liver, but it can occur in other tissues like; kidney, lungs, GIT flora, wall etc..
- The use of drugs in patients with liver disease is an art, requiring common sense, and it should be based on the evolving science of pharmacokinetics and pharmacodynamics
- Because of the liver's strategic location between the portal and systemic circulations and its extensive enzyme activity, the liver plays a vital role in drug metabolism.

Role of The Liver in Dealing with Drugs:

- Inctivation of most of the drugs.
- Activation of some drugs.
- Detoxification of most of the drugs.
- Production of more toxic metabolites of some drugs.
- Synthesis of plasma proteins mainly albumin, with its well known
 Drug binding ability.
- Secretion of drugs with bile, with consequent enterohepatic recycling.
- The function of the liver is to transform lipophilic substances into hydrophilic ones to be eliminated in urine or bile, or other aquous media.
- This is achieved by two types of reactions:

Phase I: it includes: oxidation, reduction, and hydrolysis reactions.

Aiming at introduction of a functional group to render the molecule polar

Phase II: includes conjugation reactions with: glucuronic acid, acetyl Co-A, glutathione, glycine, phosphoadenosy l-phosphosulfate (sulphate congugation), S- Adenosyl Methionine (Methylation).

N.B. The two phases are not sequential, and any of them may precede or come after the other.

- Drug biotransformation is achieved by the cytochromes
- (Microsomal and non-microsomal enzyme systems).

- More than 100 members of these cytochromes are known and identified. They are involved in drug metabolism.
- They are members of four gene families; (P450-I, II, III, IV).
- Individual p450s have distinct catalytic properties, and can metabolize many drugs and have unique binding sites and have substrate specificity.
- Genetic polymorphism of enzymes can lead to altered drug handling by liver, e.g. Slow and fast acetylation of INH due to polymorphism of acetylation enzyme.
- Genetic difference in their catalytic properties may lead to idiosyncratic response to drugs.
- Abnormal reactors for drug metabolism can be identified by PCR amplification of mutant genes.

Factors Affecting Bio-Transformation of Drugs

- 1. Inducers and inhibitors of the CYP, MES.
- 2. Hepatic blood flow; esp. For drugs with high hepatic first pass metabolism.
- 3. Malnutrition.
- 4.Age: young age:decreased due to deficint enzyme synthesis
 - Old age: aging enzymes, decrease Hepatic BI.F. & Hepatic volume.
- 5.Sex: more activity in males than females.
- 6.Liver diseases; impaired metabolism is proportionate to HC affection. It decreases clearance of some but not all drugs.
- 7. Genetic factors
- Drug biotransformation is impaired in presence of liver diseases. In patients with acute or chronic liver disease, drugs depressing CNS may precipitate encephalopathy.
- Glucuronidation is relatively spared. Hence degradation of drugs by conjugation is less affected. This may be because there are extra hepatic glucuronidation pathways or the glucuronyl transferases may be so located in the interior of microsomes that they are protected from injury.
- Benzodiazepines such as diazepam and

- chlordiazepoxide undergo oxidative biotransformation and hence clearance is impaired in liver disease, whereas oxazepam and lorazepam are inactivated by glucuronidation and hence clearance may not be affected in stable cirrhotics.
- Hepatic acetylation is also impaired in patients with liver disease. INH acetylation is hence impaired.
- No reliable marker of hepatic drug metabolizing function that predict changes in hepatic clearance, analogous to CrCI in renal drug clearance. PT & S. Albumin may help.

Mechanisms of Impaired Drug Disposition in Liver Disease

- 1. Reduced enzyme activity
- 2. Parenchymal cell mass reduced
- 3.Reduced sinusoidal perfusion and capillarisation of sinusoids
- 4. Porto systemic shunting
- 5. Impaired bile flow
- 6. Changes in protein binding
- 7. Associated decreased renal clearance, malnutrition, Endocrine changes.
 - Hepatic clearance of drugs depends upon:
 - a. Efficiency of drug metabolizing enzymes.
 - b. Hepatic blood flow(hepatic perfusion).
 - c. Intrinsic clearance(extraction) of drugs by the liver.
 - D. Extent of protein binding of drugs.
- Drug bioavailability is also controlled by the liver's capacity to clear the drug from the circulation.
- This depends on both hepatic blood flow and the efficiency of drug removal by hepatocytes (extraction ratio).
- If the latter is very high, drug clearance primarily depends on hepatic blood flow (e.g., Propranolol, Lidocaine), whereas flow has relatively little effect on drugs that are slowly cleared by the liver (e.g., Theophylline, Warfarin, Diazepam).

- Most drugs are cleared at intermediate rates, which are affected by alterations in both hepatic flow and extraction capacity.
- $CL_H = QH \{C_A CV / C_A\} = QE$ (extraction ratio)

[Q is the hepatic blood flow = 90 l/hr]

- If hepatic Cl>hepatic blood flow, Bioavailability is low, and v.v.
- If hepatic blood flow is decreased e.g. CHF, Cirrhosis, Beta Blockers, Cimetidine, etc... there will be increased systemic effects of Hepatic first pass drugs; e.g. Lidocaine, verapamil, propranolol, INH, Morphine, several Tricyclic Anti-Depressants.

CYP₄₅₀ Inducers:

- They increase synthesis of cyp enzymes. or:decrease degradation of these enzymes, e.g.Troleandomycin.
- Examples: Barbiturates except Secobarbital, Rifampicin, Phenytoin, Phenylbutazone, Griseofulvin, Smoking, Troleandomycin, Insecticides, Hydrocarbons, Androgens.
- Ethyl alcohol in small dose is enzyme inducer, meanwhile, it is enzyme inhibitor in large doses.
- Omeprazole (Proto Pump Inhibitor, used in treatment of peptic ulcer), is an inducer of CYP 1A_{1&2}, meanwhile it is an inhibitor of other CYPs.
- The effect of enzyme inducers is slow, unlike enzyme inhibitors whose effect is rapid.

Clinical signiffcance of enzyme inducers:

- Decrease the response to the inducer itself. (This may explain tolerance to some drugs e.g. Barbiturates).
- Decrease the bioavailability and hence the response to concomittant drugs e.g. Oral antidiabetics (leading to hyperglycemia), oral anti-coagulants(leading to thrombosis) & oral contraceptives(leading to pregnancy and break-through bleeding.
- Increase toxicity of metabolite induced tissuetoxicity e.g. INH, Paracetamol.

CYP₄₅₀ Inhibitors

• Examples: Allopurinol, Chloramphenicol, Isoniazid, Cimetidine, Dicumarol, Disulfiram,

Ethanol, Ketoconazole, Nortriptyline, Oral Contraceptives, Secobarbital, Troleandomycin, Macrolides, Quinolones, Anti-histaminics. Bio-flavonoids of grape-fruits juice. ---> **They act through:**

- Competitive inhibition. (Cimetidine, Ketoconazol)
- Catalytic Inhibition through comlex formation (Troleandomycin)
- Alkylation of the protein of the enzyme (Allopurinol, INH)
- Alkylation of the heme of the enzyme (Oral Contraceptives, Ethanol).
- Alkylation of the heme and protein of the enzyme (secobarbital).

Clinical signiff cance of enzyme inhibitors:

- o Increase bioavailability and henceeffect of concomittant drugs e.g. Oral anticoagulants(leading to bleeding) oral hypoglycemics (leading to hypoglycemia).
- o Increase toxicity of Concomittant Drugs, e.g. Grape fruit juice causes the arrhythmias if combined with Macrolides or Quinolones.
- Decrease metabolite induced toxicity, e.g. Cimetidine reduces toxicity of acetaminophen.

Genetic determinants of drug metabolism and drug interactions

There are several enzymes that are responsible for the hepatic metabolism of drugs. There is inherited variability in their concentration and metabolic activity (Polymorphic Metabolism): N-Acetyltransferase N-Acetyltransferase (NAT) is responsible for the transformation of procainamide to its active metabolite N-Acetylprocainamide (NAPA) and for the metabolism of other drugs such as isoniasid and hydralazine. About 50% of Caucasians and African-Americans are "Slow' Acetylators, a Phenotype that is rare among Asians. This pattern arises because of the existence of two NAT isoforms which are products of different genes. Nat₁ is expressed in virtually all individuals, while NAT₂ is expressed only in rapid acetylators.

Slow Acetylators of Procainamide have higher parent drug concentrations and lower NAPA concentrations that do fast Acetylators. This is probably not important for antiarrhythmic activity, since NAPA exerts some electrophysiologic actions of its own which are different from those of procainamide (i.e., Class iii versus class ia activity). Drug-induced lupus accumulation of procainamide may have deleterious effects due both to the native drug and to subsequent metabolism to toxic metabolites such procainamide hydroxylamine. This explains why slow acetylators of procainamide are at greater risk for the development of drug-induced lupus during procainamide therapy than rapid acetylators. Similar considerations apply to isoniazid and hydralazine. In comparison, there is little or no antinuclear antibody formation or symptomatic lupus following the administration of NAPA.

Cytochrome P₄₅₀:

Historically, cytochrome P_{450} has been conceptualized as one enzyme or an enzyme complex with the capacity to metabolize a number of drugs by a variety of routes.

More recent information has shown that there are over 100 isoenzymes of Cytochrome P_{450} , and the capacity of these enzymes is more clearly delineated with current research.

Nomenclature for the P_{450} enzymes is based upon amino acid sequence Homology.

Enzymes that have 40% or more homology are assigned the same arabic numerical, those with 55% or more homology are assigned the same letter. As an example, the sterol 27-hydroxylase and vitamin D3 24 hydroxylase enzyme are more than 40%, but less than 55% homologous, and are assigned to the CYP $_{27}$ family as CYP $_{27a}$ and CYP $_{27b}$ respectively.

The importance of characterizing the functions of different CYP (P_{450}) isoenzymes is twofold:

- First, it allows studies of population of patients to discern different phenotypes of drug metabolism.
- The second role of characterization of substrate specificity of cytochrome p450 isoenzymes is its importance in drug interactions. In particular, specific inhibitors of some isoenzymes have been identified.

CYP_{2d6} Isoenzyme

- This is one of the best-characterized ${\rm CYP}_{_{450}}$ isoenzymes.
 - It metabolizes a variety of drugs as will be discussed subsequently.
 - There are clear phenotypes of this enzyme, and there are racial differences in this expression. as an example, up to 8-10% of Caucasians are deficient in this enzyme and are called poor metabolizers of those drugs metabolized through this mechanism.
 - In contrast, the Asian populations have a very low incidence of poor metabolizers (≥ 1%).

Multiple molecular mechanisms have been identified in subjects who are poor metabolizers, resulting in the production of reduced amounts of nonfunctional CYP_{2d6} protein. In such "poor metabolizers", the functional consequences of enzyme deficiency depend upon the presence of other routes of drug metabolism and on the activity of any metabolites whose formation requires functional CYP_{2d6}. On the other hand, individuals have been described in whom two or more copies of the CYP_{2d6} gene are present. CYP_{2d6} activity may be exaggerated in these "hyperextensive metabolizers".

- CYP_{2d6} is a major route of biotransformation of a number of drugs including:
 - o Antiarrhythmic agents: Propafenone, Flecainide
 - o Beta-Blockers: Timolol, Propranolol, Metoprolol, Labetalol.
 - o Antidepressants: SSRIS as Fluoxetine, Paroxetine, Fluvoxamine, and some Tricyclic Antidepressants as Imipramine, Desipramine, Amitriptyline and Nortriptyline.
 - Statins- Simvastatin and some other members of this class, analgesics such as Codeine.

The importance of the CYP_{2d6} polymorphism vary with the different drugs:

o Increased drug activity with β -blockers, the CYP_{2d6}-dependent metabolites are less active than the parent drugs. As a result, poor metabolizers, who have higher concentrations of the parent

- drug, display a greater degree of β -blockade than extensive metabolizers. Similarly, propafenone, but not its metabolites, exerts β -blocking activity in vitro. Thus, at any given dose, the plasma propafenone concentration and therefore the degree of β -blockade are greater in poor compared to extensive metabolizers. The importance of CYP_{2d6} phenotype is also seen with Simvastatin as it can affect both the degree of lipid lowering and tolerability.
- o Little change in drug activity the clinical significance of CYP_{2d6} is less with Flecainide. Although Flecainide biotransformation to inactive metabolites is CYP_{2d6}-dependent, the unchanged parent drug can undergo renal excretion. As a result, the CYP_{2d6} polymorphism is ordinarily unimportant during Flecainide therapy, except in subjects with renal failure. In this setting, marked Flecainide accumulation and potential cardiotoxicity may occur in poor metabolizers.
- o Reduced drug activity Codeine undergoes CYP_{2d6}-dependent biotransformation to a more active metabolite, Morphine. Thus, Codeine has less pharmacological activity in poor metabolizers than extensive metabolizers.
- Drug inhibition of enzyme activity a number of drugs are potent inhibitors of CYP₂₄₆, including ultra low dose of Quinidine (e.g. 50mg) and some ssris (Paroxetine. Fluoxetine). Extensive metabolizers receiving these drugs will display the phenomenon of "Phenocopying" behaving as if CYP_{2d6} activity were absent. This can be illustrated by the following observations: blockade of CYP_{2d6} by ssris can increase the incidence of side effects associated with Tricyclic Antidepressants. Blockade of CYP_{2d6} by Quinidine can increase the degree of β-blockade in extensive metabolizers receiving Timolol eye drops and in those treated with Propranolol or Propafenone.

These drug interactions are genetically determined, occurring only in extensive metabolizers in whom functional CYP_{2d6} is present

66

to inhibit. Inhibition of drug activity can also occur competitively when two drugs metabolized by CYP_{2d6} are given concurrently. In one study (holtzman, et. al, 1987), healthy subjects were given both flecainide and propranolol; the plasma levels were increased by 20% for flecainide and 30% for propranolol.

CYP_{3a4} isoenzyme

• It is the most important hepatic enzyme for drug metabolism. It is responsible for the biotransformation of many commonly used drugs such as quinidine, lidocaine, statins, calcium channel blockers, theophylline and cyclosporine. It has received considerable attention, largely because this isoenzy meisthe major route of metabolism for cyclosporine as well as for other frequently used drugs, which can therefore inhibit the metabolism of cyclosporine. In contrast to CYP_{2d6}, genetically determined "poor metabolizers" with absent or low CYP3a4 activity have not been identified. However, the activity of the enzyme varies considerably among individuals for reasons that are incompletely understood.

Ingestion of a Number of Drugs Can Affect CYP_{3a4} Activity:

- Enzyme activity is reduced by erythromycin, clarithromycin, ketoconazole, itraconazole and large quantities of grapefruit. Grapefruit juice or unprocessed grapefruit appears to act by down regulating intestinal CYP₂₀₄.
- Enzyme activity is increased by rifampicin, anticonvulsants such as phenytoin and phenobarbital, and dexamethazone. These drugs appear to activate the nuclear pregnane X receptor (PXR), which binds to a response element in the CYP_{3,34} promoter. The ability of CYP_{3a4} to be induced and its broad substrate specificity are responsible for its drug interactions. It is also possible that endogenous substances modulate the expression or function of CYP_{3a4}, resulting in a wide variation in enzyme activity. Some of the interindividual variability may be genetically determined.

Competitive inhibition of CYP_{3a4} can lead to clinically important drug interactions. The effects are most pronounced when there is no alternative

pathway for the metabolism of a potentially toxic drug. The most spectacular example is the development of torsade de pointes during the administration of terfenadine or cisapride and a potent inhibitor of CYP_{3a4}. These drugs block cardiac potassium channels and normally undergo extensive presystemic biotransformation to metabolites that have no electrophysiologic activity. The ingestion of CYP_{3a4} inhibitors (e.g. Erythromycin or Ketoconazole) or the presence of a drug overdose or liver disease results in the accumulation of the native drug, resulting in qt prolongation and the potential for torsade de pointes.

The use of a potent CYP_{3a4} inhibitor, especially cyclosporine, can increase the plasma concentrations of simvastatin and most other statins, enhancing the risk of myopathy. Weaker inhibitors, such as the ccbs, have less risk. A similar type of drug interaction explains the increased cyclosporine concentration seen with co-administration of Ketoconazole, Diltiazem, or grapefruit juice. On the other hand, drugs that increase enzyme activity (e.g. Anticonvulsants) can reduce cyclosporine concentrations, requiring an increase in the cyclosporine dose.

CYP_{2c} Isoenzyme

This isoenzyme metabolizes the following drugs: -anticonvulsants: phenytoin, mephenytoin, methobarbital. -Antimalarials: proguanil, cycloguanil -oral hypoglycemic: tolbutamide -anticoagulants: warfarin CYP_{2c} is absent in about 3% Of caucasian patients, and in substantially higher numbers of asian (greater than 10%). Cimetidine inhibits this isoenzyme, although it may also have effects on other cytochrome p450 isoenzymes. Thus, the administration of cimetidine with one of the above drugs may lead to a clinically important drug interaction.

CYP_{2c9} Isoenzyme

This isoenzyme inactivates the S-enantiomer (mostactive) of warfarin. There are multiple genetic polymorphisms of CYP_{2c9}. The maintenance doses required to reach a target inr can vary according to the genotype. Amiodarone inhibits CYP_{2c9}. This explains the potentiation of warfarin effect by amiodarone.

CYP_{2c19} Isoenzyme

Omeprazole, lansoprazole, and to a variable

The activity of CYP_{2c19} affects the likelihood of success in the treatment of both helicobacter pylori and gastroesophageal reflux disease (GERD):

• The effect of variable metabolism of omeprazole in the treatment of helicobacter pylori was examined in japanese patients (Furuta, et. al,1998). While eradication was achieved in all individuals homozygous for a CYP_{2c19} mutation, successful treatment was achieved in only 60 and 29% of heterozygotes and wild type homozygotes, respectively. Another report from japan (Furuta, et. al, 2002) evaluated the efficacy of lansoprazole in the treatment of gerd. Poor metabolizers were much more likely to be cured on endoscopic examination than heterozygotes and wild type homozygotes (85 versus 68 and 46 % respectively). The cure rate was only 16% in wild type homozygotes who had severe GERD. Wild type homozygotes had the lowest plasma lansoprazole concentrations.

The CYP_{2c19} genotype also affects the metabolism of sertraline. Plasma sertraline concentrations are significantly elevated in patients with a CYP_{2c19} mutation. The clinical significance of this difference is uncertain.

N.B: a drug like Rifampicin induces a variety of hepatic enzymes, and, within the Cytochrome P_{450} isoenzymes, produces a variable degree of induction. In an in vitro study of human hepatocytes (Rae, et al, 2001), Rifampicin substantially induced the CYP_{3a} family, CYP_{2c9}, CYP_{2a6}, and a number of non-P₄₅₀ enzymes but had little effect on CYP_{2d6} or CYP_{2e1}. Further research will undoubtedly identify other isoenzymes of Cytochrome P₄₅₀ and will refine our knowledge as to specific substrates or inhibitors. This will allow more precise prediction of clinically important

drug interactions in humans.

Pseudocholinesterase Deffciency

The paralytic agent succinylcholine is metabolized by the circulating enzyme pseudocholinesterase. The enzyme is functionally absent in approximately 1 in 3000 individuals. In such patients, prolonged paralysis may occur following ordinary doses of succinylcholine.

Thiopurine Methyltransferase

methyltransferase Thiopurine (TPMT) is responsible for the biotransformation of antimetabolites such as 6-mercaptipurine and azathioprine to less active metabolites. Rare cases of mutations in the tpmt gene have been described; during treatment with one of the above drugs, affectedsubjectsdisplayprolongedagranulocytosis that can be fatal or develop lower leukocyte counts necessitating drug discontinuation. In one series, for example 6 of 67 patients receiving azathioprine for rheumatic disease where heterozygous for mutant tmpt alleles (Black et. al., 1998). Five of these patients had to discontinue therapy within one month because of leucopenia, while the sixth did not adhere to therapy. In comparison the median duration of therapy was 39 weeks in those with wild type alleles. Some data also suggest that increased activity of tpmt may be associated with a decreased antitumor efficacy of thiopurine drugs.

Similar considerations apply to the use of 6-mercaptopurine. In a series of 180 children with acute lymphoblastic leukemia, the cumulative incidence of dose reductions in 6-mercaptopurine due to drug toxicity was 7% in the 161 children with two wild-type alleles, 35% in the 17 children with one mutant allele, and 100% in the two children with two mutant allele. Lowering the 6-mercaptopurine dose in the last two groups permitted administration of the full chemotherapy protocol.

P-glycoprotein

p-glycoprotein ($P_{\rm gp}$), also called multidrug resistance glycoprotein, is adrug transport system that was initially identified in certain tumor cells that displayed the phenomenon of drug resistance because the cells were able to pump various anticancer drugs out of their cytoplasm. In addition to tumor cells, $P_{\rm gp}$ is expressed on the luminal aspect of enterocyes, on the canalicular aspect of hepatocyes, the luminal aspect of renal tubular cells, and in the capillaries of the blood-

brain barrier. Substrates for $P_{\rm gp}$ include anticancer drugs, digoxin, quinidine, cyclosporine, and many others

 $\boldsymbol{P}_{\text{gp}}$ expression may have a number of clinically important effects:

- Animal and in vitro studies suggest that poor central nervous system penetration of hiv protease inhibitors may be due to P_{gp} mediated efflux at the capillaries of the blood-brain barrier.
- The elevation in serum digoxin induced by a variety of drugs (quinidine, verapamil, cyclosporine, erythromycin, ketoconazole) is probably due to inhibition of P_{gp} mediated digoxin transport; especially since digoxin does not undergo significant metabolism prior to excretion. The administration of agents that inhibit P_{gp} by reverting the multidrug resistance phenotype (e.g. Quinine), by inhibiting the activity and maturation of P_{gp} (e.g. Cyclosporine), may increase the response to chemotherapeutic agents.

Pharmacokinetic and Pharmacodynamic

Changes in Drug Disposition in Hepatic Impairment

Absorption:

- It is impaired from GIT due to several factors:
 - o. Nausea and vomiting with hepatic or biliary dysfunction.
 - o Dyspepsia and indigestion which interfere with drug absorption.
 - o Non-healthy mucosa of GIT.
 - o Intestinal mucosal congestion in cases of portal Hypertension&chf.
 - o Defective biliary secretion with consequent pancreatic enzyme

 Dysfunction, which interfere with activation of some drugs.
 - o Use of polypharmacy with possible hazardous interactions, or interference with intestinal flora.

• Protein Binding:

- Liver is the site of synthesis of plasma proteins,particularly Albumin. Thus with liver impairment,there will be hypoabuminemia, and free parts of drugs will increase, needing dose adjustment.

• Distribution:

- o In cases of portal hypertension with ascitis, and in cases of generalized oedema with liver cirrhosis, the volume of distribution is increased especially for hydrophilic drugs, which necessitates dose adjustment.
- o Decreased hepatic blood flow with CHF, and liver cirrhosis, will increase the systemic bioavailability of drugs known to undergo first pass hepatic metabolism.

• Biotransformation:

- o Impairment of biotransformation of drugs metabolized in the liver.
 - (This point is detailed beforehand).
- o Increased susceptibility of the liver to drug insults.

• Elimination:

- o Decreased biliary elimination of drugs eliminated by this route.
- o Co-morbidity of the other organs of elimination e.g. Kidneys, may occur.

• Organ sensitivity:

o Hepatic patients are more susceptible to CNS depressants.

Effect of Liver Disease on Some Pharmocologic Agents

- **1.** Tranqulizers and sedatives: alprazolam, diazepam, chlordiazepoxide, midazolam-half life is increased by 50% to 400 % and clearance reduced by 30%-60%. Barbiturates if albumin levels normal nodosage adjustment required
- **2.** Antidepressants and anticonvulsants:

 *phenytoin no change in drug disposition
 in acute viral hepatitis. *Amitryptiline,
 fluoxetine, valproic acid half life is increased
 and clearance decreased

3. Analgesics and NSAIDS:

- **Ibuprofen and phenyl butazone** no change in drug disposition dextropropoxyphene and pentazocine-clearance is impaired phenacetain-caution in use in alcoholic liver disease
- **4.Cortico Steroids:** no change in drug disposition

5.Cardio Vascular Drugs:

- Antiarrythmics clearance is decreased and half life increased.
- Beta blockers: labetolol and metoprolol no change -propranalol clearance is decreased, half life increased calcium channel blockers clearance is decreased half life increased
- Anti hypertensives ACE inhibitors, NO change, diuretics half life and clearance altered especially in presence of renal failure. Digoxin, NO change
- **6.Gastric Acid Suppressants:** only proton pump inhibitors have low clearance and increased half life.
- **7.Xanthines:** Theophylline, clearance decreased with increased half life.
- **8.Antibiotics:** Chloramphenicol, Clindamycin, Vancomycin, Erythromycin, Cefoperazone, Rifampicin, Half life is increased consequent to decreased clearance
- **<u>9.Anticoagulants:</u>** clearance unchanged but bioavailability increased
- **10. Oral Hypoglycemics:** Hypolipidemics such Gemfibrosil, Colchicine, Metoclopramide, Bioavailability is increased.

Note

- Drug utilization in hepatic patients:
- Prodrugs of ACEI, are better avoided. Use Captopril or lisinopril.
- Antacids: avoid sodium containing prep. In ascites. Avoid Alum. Prep. Which may precipitate coma due to constipation.
- Azithromycin, Clarithromycin are better avoided for fear of jaundice (common).
- Cimetidine increases risk of confusion.
- Dantrolene causes severe liver damage.
- Reduce dose of Cal. Channel Blockers.
- Ketotifen increases risk of sedation.
- Reduce dose of Metronidazole, give it once/day in liver diseases.
- Neomycin is absorbed in liver disease from GIT, with increased risk of Ototoxicity.
- Pyrazinamide may lead to idiosyncr. Hepatic toxicity in liver diseases.
- Quinolones:avoid Ciprofloxacin,no Rfloxacin, Nalidixic acid.
 Reduce dose of Ofloxacin.
- Valproate: better avoided in liver diseases.

Guideline Development Group Acknowledgements

1 - Cairo University Consultancy group:

Professor Dr. Laila Kamel	Professor of Public Health& Community Medicine Cairo University, Faculty of Medicine
Professor Dr. Nagwa Eid	Professor of Internal Medicine, Cairo University, Faculty of Medicine
Professor Dr. Salma Dawara	Professor of General Surgery Cairo University, Faculty of Medicine
Dr. Abeer Barakat	Lecturer of Public Health Cairo University, Faculty of Medicine

2- Sector of Technical Support and Project Technical Working & Supervisory Group:

Dr. Emam Moussa	Head of the Central Administration of the Technical Support and Project & Group Leader
Dr. Soad Abdel Megid	Guideline Developer & Groups Coordinator
Dr. Osama Abdel Azim	Technical Advisor

3- Additional Support and First Draft Revision:

MOHP Level

All 1st Undersecretary , and Undersecretary of the MOHP Sectors and Central Administrations are involved in revising the Document

Medical University Staff and Institutions		
Professor Dr. Mahmoud Serry	Professor of Chest, Ein Shams University	
Professor Dr. Omima El Gebally	Professor of Family Med, Assiut University	
Professor Dr. Fathy Maklady	Professor of Family Med & General Medicine, Canal El Suez University	
Professor Dr. Esmat Shiba	Professor of Family Med & General Medicine, Cairo University	
Professor Dr. Hesham Zaher	Professor of Dermatology, Cairo University	
Professor Dr. Ezz El Dine Osman	Professor of OB/Gyn, Cairo University	
Professor Dr. Tarek Kamel	Professor of ENT, Cairo University	

Professor Dr. Magda Badawy	Professor of Pediatric, Cairo University
Professor Dr. Tagrieed Farahatt	Professor & Head Department of Family medicine, Monofiya University
Professor Dr. Sawsan Fahmy	Professor of Public Health, Alex, High Institute of Public Health
Professor Dr. Osman Ziko	Professor of Ophthalmology, Ein Shams University
Professor Dr. Amr El Noury	Consultant of Clinical Guideline MOHP General Hospital

4- Revision of Pharmaceutical Sections :

Professor Dr. Hider Galeb	Professor of Pharmacology, Cairo University
Professor Dr. Abdel Rahman El Nagar	Professor of Clinical Pharmacology, Faculty of Medicine, Cairo University
Professor Dr. Aza Monier Agha	Professor of Pharmacology, Faculty of Pharmacy, Cairo University
Professor Dr. Faten Abdel Fatah	Professor of Clinical Lab " Institute of Pharmaceutical Monitoring"
Dr. Mohamed Awad	Lecturar of Pharmacology, Faculty of Pharmacy, Helwan University
Dr. Alaa Mokhtar	Director of HSRP Pharmaceutical Program
Dr. Gebriel Ali	MOHP, Information Center, Central Administration of pharmacy
Dr. Mostafa Sleim	Pharmaceuticals Consultant at MOHP

5- Revision By High Committee of Egyptian Board of Family Medicine

Professor Dr. Gabr Metwally	Professor of Public Health Al Azher Unversity for boys
Professor Dr. Mohamed Farghally	Professor of Public Health Al Azher Unversity for boys
Professor Dr. Adel Fouda	Professor of Public Health Zagazig University

6- Family Physician Participating in the Review of the First Draft and Field testing of the Document at Governorate levels:

Governorate level		
"Sohag"		
Dr. Mazhar Attia Ahmed	El Shik Shebl, FHU	
Dr. Gerges Khalil Krns	El Gazazra FHU	
Dr. Emad Latif Metiass	El Shik Yosef FHU	
Dr. Komyl Wdiee Danial	Bahta FHU	
Dr. Kadry Mohamed Attia	Erabat Abu Ezize FHU	
Dr. Emad Naeeim Loka	Bahatyl El Gizira FHU	
Dr. Hala Samuaeil Fares	TST Quality Specialest	
Dr. Frag Ahmed Mahmoud	TST Primary Health Care Director	
Governorate level '' Qena''		
Dr. Nahla Shikhoon	El Mkrbya FHU	
Dr. Nesreen Abu El Abass Elian	El Hragia FHU	
Dr. Eiman Mohamed Mahfouz	Gzyra Motira FHU	
Dr. Mona Fakhry Ali	El Hogirat FHU	
Dr. Mohamed Mohamed Ashour	El Homer Wal Gaafraa FHU	
Dr. Mostafa Glal Osman	El Tob FHU	
Dr. Ahmed Saad Ahmed	TST Coordinator	
Dr. Mamdouh Abuel Kasem	TST	
Governorate level '' Monoffya''		
Dr. Tamer Farag Ali	Mastay FHU	
Dr. Alaa El Dine Abdel Razek	Ashliem FHU	
Dr. Sherif Mosaad Labib	El Remally FHU	
Dr. Asmaa Mahmoud El Sayed	Shobra Bakhom FHU	
Dr. Waleid Mohamed Rashad	Meit Bara FHU	
Dr. Nahed Sobhy Mahmoud	Tymor FHU	
Dr. Gehad Ibrahim Mohamed	TST	

Governorate level '' Alexandria''		
Dr. Naira Niazy	Alexandria Central Coordinator	
Dr. Nagwa Mostafa Abuel Nazar	El Gomrok FHU	
Dr. Ghada Mohamed Abdel Allah	El Gomrok FHU	
Dr. Marian Nashaat	El Manshia2	
Dr. Ihab Zaky Iraheeim	El Laban1 FHU	
Dr. Riham Sabry	El Laban1 FHU	
Dr. Nadia Khaliel Fahmy	El Laban2 FHU	
Dr. Anas Mohamed Helal	TST	
Dr. Maha Mogib Haseib	TST	
Dr. Sanaa Ahmed Elbarsiky	TST	
Governorate level ''Suez''		
Dr. Zein El Abedein Abdel Motelb	El Safaa FHU	
Dr. Saher Mahmoud Hussien	El Amal FHU	
Dr. Amany Keshk	El Sweiz1 FHU	
Dr. Mervet Gharieeb	El Mothalath FHU	
Dr. Suzan Gamiel	24 October FHU	
Dr. Hany Anter	El Mashroo FHU	
Dr. Nadia Mohamed Esmaeil	TST	
Dr. Magda Ahmed Mohamed	TST Coordinator	